



Primary Fungal Prophylaxis in Hematological Malignancy: a Network Meta-Analysis of Randomized Controlled Trials

 Cho-Hao Lee,^a Chin Lin,^{b,c} Ching-Liang Ho,^a Jung-Chung Lin^d

^aDivision of Hematology and Oncology Medicine, Department of Internal Medicine, Tri-Service General Hospital, National Defense Medical Center, Taipei, Taiwan, Republic of China

^bSchool of Public Health, National Defense Medical Center, Taipei, Taiwan, Republic of China

^cDepartment of Research and Development, National Defense Medical Center, Taipei, Taiwan, Republic of China

^dDivision of Infectious Diseases and Tropical Medicine, Department of Internal Medicine, Tri-Service General Hospital, National Defense Medical Center, Taipei, Taiwan, Republic of China

ABSTRACT Several new antifungal agents have become available for primary fungal prophylaxis of neutropenia fever in hematological malignancy patients. Our aim was to synthesize all evidence on efficacy and enable an integrated comparison of all current treatments. We performed a systematic literature review to identify all publicly available evidence from randomized controlled trials (RCT). We searched Embase, PubMed, the Cochrane Central Register of Controlled Clinical Trials, and the www.ClinicalTrials.gov website. In total, 54 RCTs were identified, including 13 treatment options. The evidence was synthesized using a network meta-analysis. Relative risk (RR) was adopted. Posaconazole was ranked highest in effectiveness for primary prophylaxis, being the most favorable in terms of (i) the RR for reduction of invasive fungal infection (0.19; 95% confidence interval [CI], 0.11 to 0.36) and (ii) the probability of being the best option (94% of the cumulative ranking). Posaconazole also demonstrated its efficacy in preventing invasive aspergillosis and proven fungal infections, with RR of 0.13 (CI, 0.03 to 0.65) and 0.14 (CI, 0.05 to 0.38), respectively. However, there was no significant difference among all of the antifungal agents in all-cause mortality and overall adverse events. Our network meta-analysis provided an integrated overview of the relative efficacy of all available treatment options for primary fungal prophylaxis for neutropenic fever in hematological malignancy patients under myelosuppressive chemotherapy or hematopoietic cell transplantation. On the basis of this analysis, posaconazole seems to be the most effective prophylaxis option until additional data from head-to-head randomized controlled trials become available.

KEYWORDS antifungal agents, antifungal therapy, hematology, meta-analysis

Invasive fungal infections (IFIs) are a significant cause of morbidity and mortality following dose-intensive chemotherapy or hematopoietic cell transplantation in patients with neutropenic fever. The risk of IFIs is particularly increased in hematological malignancy patients (1, 2). Furthermore, invasive mold infections often occur exclusively in high-risk patients with profound neutropenia (<100 cells/mm³) that lasts longer than 10 to 15 days (3–5). Now that the threats posed by bacterial and viral infections have been somewhat reduced, IFIs have become one of the main infective causes of mortality in this population (6).

Currently, *Aspergillus* and *Candida* species account for 95% of all cases of IFIs, but the epidemiological characteristics of IFIs evolve due to the selection pressure of antimicrobials and other factors (7, 8). With the increasing use of intensive immunosuppressive cancer therapeutic modalities (9, 10), IFIs have become an important reason for delays and reduced response to therapies for hematological cancer (11–13).

Received 23 February 2018 Returned for modification 6 April 2018 Accepted 19 May 2018

Accepted manuscript posted online 4 June 2018

Citation Lee C-H, Lin C, Ho C-L, Lin J-C. 2018. Primary fungal prophylaxis in hematological malignancy: a network meta-analysis of randomized controlled trials. *Antimicrob Agents Chemother* 62:e00355-18. <https://doi.org/10.1128/AAC.00355-18>.

Copyright © 2018 American Society for Microbiology. All Rights Reserved.

Address correspondence to Jung-Chung Lin, linjungchung1@yahoo.com.tw.

Over recent decades, a series of studies have assessed the effect of antifungal agents on the prevention of IFIs. Although numerous antifungal agents are available, IFIs remain a serious problem because of obstacles to timely diagnosis and protracted initiation of therapy, the high morbidity and mortality rates associated with limited activity of antifungal agents, drug side effects, and increasing use of high-dose corticosteroids (14). Therefore, primary prevention of fungal infections, which had been repeatedly demonstrated to reduce IFIs and all-cause mortality, remains essential (15, 16). The ideal prophylactic antifungal therapy should be safe and well tolerated for long-term use, being effective against a wide spectrum of organisms and available as intravenous and oral formulations with good bioavailability (17). Recent randomized controlled trials (RCTs) (18–21) and meta-analyses (22–24) showed a marked reduction of IFIs in patients who used triazoles and echinocandins. The currently available multiple polyenes, echinocandins, and triazoles fulfill some of the requirements for an ideal prophylactic antifungal agent; however, some areas for improvement remain.

Network meta-analysis may be a more robust methodology which allows full comparisons of all relevant interventions in a single analytical model, including those which lack head-to-head comparisons (25, 26). The clinical requirement for comparison of all relevant treatments is almost impossible for current clinical trial design because of the cost and regulatory approval-driven strategies. Commonly, new interventions are compared with placebo or current standard intervention (27, 28). The synthesis analysis takes advantage of both direct (as used in the standard meta-analysis) and indirect comparisons between a number of treatments, which may strengthen the relative efficacy estimate and allow the designation of the best treatment simultaneously.

There is no single agent that will prevent all fungal infections; therefore, careful monitoring and treatment of emergent breakthrough IFIs throughout the high-risk period are essential. The aim of this study was to examine all of the evidence from qualified randomized controlled trials that have guided antifungal choices and to compare the clinical efficacy and safety of the antifungal agents for primary IFI prophylaxis in hematological malignancy patients undergoing myelosuppressive chemotherapy or hematopoietic cell transplantation.

RESULTS

The algorithm of this systematic literature review is shown in Fig. 1. A total of 331 citations were retrieved from the databases. After removing duplicates, 286 citations were screened based on the title and abstract and 203 studies were excluded from further analysis. In the second phase, 83 full texts were screened, 29 of which were excluded. A total of 54 citations were included for qualitative analysis. These citations comprised 53 full publications and 1 doctoral dissertation (19). There were 18 double-blind trials and 23 multicenter studies. All studies had an acceptable quality assessment of trials and none exhibited high risk of bias in randomization, and allocation concealment domains were found. Thirty-three trials were open-label design; 27 trials had a quality assessment score of ≥ 5 . Overall, 12,832 cases were enrolled. The details of the search process are demonstrated in the appendix posted at <https://goo.gl/6AAXgq>, pages 2 to 6.

In total, 54 trials were identified, including 13 arms: (i) oral polyene, (ii) intravenous conventional amphotericin B (iAMB), (iii) aerosolized amphotericin B (aAMB), (iv) liposomal amphotericin B (LAMB), (v) amphotericin B lipid complex (ABLC), (vi) ketoconazole (KTCZ), (vii) fluconazole (FLCZ), (viii) itraconazole (ITCZ), (ix) voriconazole (VOCZ), (x) posaconazole, (xi) micafungin (MCFG), (xii) caspofungin (CASP), and (xiii) placebo.

Table 1 lists the summary of the characteristics of the included trials. The median age ranged from 6.08 to 62.17 years; half of the patients received intensive chemotherapy, and 27% received hematopoietic cell transplantation. The trials were distributed across North America, South America, Europe, Asia, and Africa. Most of the studies enrolled patients with an absolute neutrophil count of less than 1,000/ μ l. However, the time point for primary fungal prophylaxis administration was inconsistent among the trials.




FIG 1 PRISMA flowchart of network meta-analysis.

Network meta-analysis. Figure 2 describes the integral network for primary fungal prophylaxis of hematological malignancy patients. To include all of the trials within one framework, we assumed that there were no differences in efficacy regardless of the dosage scheme and durations.

Invasive fungal infection in the overall population. Figure 3 presents the network meta-analysis results of the overall IFI incidence in a total of 54 studies (13 arms, 12,832 cases) that used placebo as the comparator. All treatments were sorted based on their ranking, along with their relative risk (RR) and 95% confidence intervals (CI), compared with that of the placebo. The probability scores for being the most effective treatment were also listed. Among the antifungal agents, posaconazole, liposomal amphotericin B, micafungin, itraconazole, voriconazole, aerosol amphotericin B, and fluconazole all revealed significantly lower invasive fungal infection incidence, with RR ranging from 0.19 to 0.51 compared with that of the placebo. Posaconazole was ranked highest in the prevention of invasive fungal infection (RR, 0.19; 95% CI, 0.11 to 0.36; probability [P] score, 94%).

TABLE 1 Basic characteristics of included randomized trials^a

Reference	Trial design	Country, total no. of cases	Mean age (yr)	Main therapy ^f	Leukemia (%)	Inclusion criteria ^c	Treatment arm ^e		Involved results ^d	Quality (Cochrane risk of bias)
							A	B		
Donnelly et al. (55)	RCT	UK, 36	37.38	C/T	91.60	NA	ITCZ, 400 mg, p.o., QD	AMB, 100 mg, p.o., QID	1, 2, 3	4
Palmblad et al. (56)	RCT	Sweden, 107	50.47	C/T	100	>15 years, ANC < 500, >5 days	ITCZ, 200 mg, p.o., QD	Placebo	1, 2, 3	7
Akiyama et al. (57)	OP, RCT	Japan, 130	43.32	C/T	98.40	>15 yr, ANC < 500, >10 days	FLCZ, 200 mg, p.o., QD	AMB, 800 mg, p.o., TID	1, 2	3
Philpott-Howard et al. (58)	OP, MC, RCT	International, 536	45.90	C/T, 87.5%	75.50	ANC < 500, >5 days	FLCZ, 50 mg, p.o., BID	AMB, 100 mg, p.o., QID	1, 2, 4	3
Freudenhil et al. (59)	DB, RCT	Netherlands, 92	49.50	C/T	88.00	>15 yr, ANC < 1,000, >3 days	ITCZ, 200 mg, p.o., BID	Placebo	1, 2, 3, 4	6
Winston et al. (60)	DB, MC, RCT	International, 256	43.48	C/T	100	ANC < 1,000, >7 days	FLCZ, 400 mg, p.o., QD	Placebo	1, 2, 3, 4	6
Chandrasekar et al. (61)	DB, RCT	USA, 46	38.00	NA	76.10	>13 yr, ANC < 1,000, >5 days	FLCZ, 400 mg, p.o., QD	Placebo	1, 2	5
Ellis et al. (62)	RCT	Saudi Arabia, 90	0.00	NA	NA	<18 yr, ANC < 1,000, >2 days	FLCZ, 200 mg, p.o., QD	Polyene, Clotrimazole, 10 mg, p.o., BID	1, 2	3
Ninane et al. (63)	OP, MC, RCT	International, 502	6.80	C/T	52.80	<18 yr, ANC < 1,000, >2 days	FLCZ, 3 mg/kg, p.o., QD	AMB, 25 mg/kg, p.o., QID	1, 2, 4	3
Riley et al. (64)	DB, RCT	USA, 35	38.00	SCT	60	ANC < 500	AMB, 0.1 mg/kg, i.v., QD	Placebo	1, 2, 3	7
Bodey et al. (65)	OP, RCT	USA, 77	46.47	NA	100	>16 yr, ANC < 500, >5 days	FLCZ, 400 mg, p.o./i.v., QD	iAMB, 0.5 mg/kg, i.v., QID	1, 2, 4	3
Behre et al. (66)	OP, RCT	Germany, 115	43.00	C/T	67.80	>18 years ANC < 500, >10 days	aAMB, 10 mg, iH, BID	Placebo	1, 2, 3	4
Egger et al. (67)	OP, RCT	Switzerland, 89	38.42	C/T, 67%	64.00	>14 yr, ANC < 1,000, >5 days	FLCZ, 400 mg, p.o./i.v., QD	Nystatin, p.o., QID	1, 2	6
Schaffner et al. (68)	OP, RCT	Switzerland, 151	39.50	NA	37.70	ANC < 500, >5 days	FLCZ, 400 mg, p.o., QD	Placebo	1, 2, 3	7
Annaloro et al. (69)	OP, RCT	Italy, 53	33.62	SCT	NA	NA	FLCZ, 300 mg, p.o., QD	ITCZ, 400 mg, p.o., QD	1, 2, 3	3
Kern et al. (70)	OP, MC, RCT	International, 68	48.71	C/T	100	>18 ANC < 500, >7 days	FLCZ, 400 mg, p.o., QD	AMB, 40 mg, p.o., Q4H	1, 2, 3, 4	5
Kelsey et al. (71)	DB, MC, RCT	International, 161	39.92	SCT, 83.8%	72.50	>16 yr, ANC < 1,000, >5 days	AMB, 2 mg/kg, p.o., TID	Placebo	1, 2, 3, 4	7
Menichetti et al. (72)	DB, MC, RCT	Italy, 405	44.00	C/T, 63.9%	75.60	>18 ANC < 1,000, 7 days	ITCZ, 2.5 mg/kg, p.o., BID	Placebo	1, 2, 3	5
Rotstein et al. (74)	DB, MC, RCT	Canadian, 274	46.44	C/T, 56%	56.20	>18 ANC < 500, >7 days	FLCZ, 400 mg, p.o., QD	Placebo	1, 2, 3, 4	7
Schwartz et al. (75)	OP, MC, RCT	Germany, 382	46.81	NA	83.20	>18 ANC < 500, >7 days	aAMB, 10 mg, iH, BID	Placebo	1, 2, 3	5
Young et al. (76)	DB, MC, RCT	International, 164	43.23	NA	94.50	>16 yr, ANC < 1,000, >5 days	FLCZ, 200 mg, p.o., QD	Nystatin, 6MIU, p.o., QD	1, 2, 4	7
Huilgens et al. (70)	DB, RCT	Netherlands, 202	15.15	NA	38.60	>18 yr, ANC < 500, >14 days	FLCZ, 50 mg, p.o., BID	ITCZ, 100 mg, p.o., BID	1, 2, 3	5
Morgenstern et al. (73)	MC, RCT	International, 445	44.55	C/T, 78.9%	84.50	ANC < 100, >7 days	FLCZ, 100 mg, p.o., QD	ITCZ, 2.5 mg/kg, p.o., BID	1, 2	3
Harousseau et al. (77)	DB, RCT	International, 557	48.74	C/T, 81.3%	70.70	>18 yr, ANC < 500, >14 days	ITCZ, 2.5 mg/kg, p.o., BID	AMB, 500 mg, p.o., QID	1, 2, 3, 4	7
Lavardiére et al. (78)	DB, RCT	Canada, 266	46.32	C/T, 56%	58.60	>18 yr, ANC < 500, >7 days	FLCZ, 400 mg, p.o., BID	Placebo	1, 2	5
Nucci et al. (79)	DB, MC, RCT	Brazil, 210	27.77	C/T, 85.3%	80	ANC < 1,000, >7 days	ITCZ, 100 mg, p.o., BID	Placebo	1, 2, 3	7
Wolff et al. (80)	OP, MC, RCT	USA, 355	42.55	SCT	NA	>18 yr, ANC < 500, >10 days	FLCZ, 400 mg, p.o./i.v., QD	iAMB, 0.2 mg/kg, i.v., QD	1, 3, 4	3
Boogaerts et al. (81)	OP, MC, RCT	Finland, 277	46.92	C/T	67	>16 yr, ANC < 500, >10 days	ITCZ, 100 mg, p.o., BID	AMB, 500 mg, Nystatin, 2 MIU, p.o., QID	1, 2, 3, 4	3
Koh et al. (82)	OP, RCT	Singapore, 186	29.69	SCT	66.60	ANC < 1,000, >7 days	FLCZ, 200 mg, p.o., QD	iAMB, 0.2 mg/kg, i.v., QD	1, 2, 3	3
Kaptan et al. (83)	RCT	NA, 55	35.58	C/T	100	>18 yr, ANC < 500, >7 days	ITCZ, 200 mg, p.o., BID	Placebo	1, 2, 3	3
Lass-Flörl et al. (84)	RCT	Austria, 106	43.94	C/T	78.30	>18 yr	ITCZ, 5 mg/kg, p.o., QD	AMB, 1,000 mg, p.o., TID	1, 2, 3	3
Winston et al. (4)	OP, MC, RCT	International, 138	39.54	SCT	69.50	>13 yr	FLCZ, 400 mg, p.o./i.v., QD	ITCZ, 200 mg, p.o., BID	1, 2, 3, 4	4
Marr et al. (85)	OP, RCT	USA, 299	0.00	SCT	NA	>13 yr, ANC < 500, >10 days	FLCZ, 400 mg, p.o./i.v., QD	ITCZ, 2.5 mg/kg, p.o./i.v., TID	1, 2, 3	3
Van Burik et al. (86)	DB, MC, RCT	USA, 830	42.52	SCT	29.20	≥6 months, ANC < 500, >4 days	FLCZ, 8 mg/kg, i.v., QD	MCFG, 1 mg/kg, i.v., QD	1, 2, 3, 4	7
Hiemenz et al. (87)	DB, RCT	USA, 74	43.24	SCT	NA	≥18 yr, ANC < 500	FLCZ, 400 mg, i.v., QD	MCFG, 5 mg/kg, i.v., BID	1, 2, 3, 4	3
Penack et al. (91)	OP, RCT	Germany, 219	53.76	NA	79.50	>18 yr, ANC < 500	LAMB, 50 mg, i.v., QD	Placebo	1, 2, 3	5
Glasmacher et al. (88)	OP, RCT	Finland, 494	0.00	NA	88.60	ANC < 500, >10 days	FLCZ, 400 mg, p.o./i.v., QD	ITCZ, 5 mg/kg, p.o., BID	1, 2, 3, 4	5
Oren et al. (90)	RCT	Israel, 195	49.49	SCT	43.50	>15 yr, ANC < 500, >7 days	FLCZ, 400 mg, p.o./i.v., QD	ITCZ, 200 mg, p.o./i.v., BID	1, 2, 3	3
Mattuzzi et al. (89)	OP, RCT	USA, 197	62.17	C/T, 93.4%	75.10	>15 yr, ANC < 500, >7 days	ITCZ, 200 mg, i.v., QD	CASP, 50 mg, i.v., QD	1	5
Vehreschild et al. (93)	DB, MC, RCT	Germany, 25	53.60	NA	NA	>18 yr, ANC < 500, >5 days	VOZC, 200 mg, p.o., BID	Placebo	1, 2, 4	7
Ullmann et al. (34)	DB, MC, RCT	International, 579	41.30	SCT	71.8	>13 yr, ANC < 500, >7 days	FLCZ, 200 mg, p.o., TID	p.o.CZ, 200 mg, p.o., TID	1, 2, 4	5
Cornely et al. ^b (27)	OP, MC, RCT	International, 592	49.44	C/T	86.20	>13 yr, ANC < 500, >7 days	p.o.CZ, 200 mg, p.o., TID	FLCZ, 400 mg, p.o., QD	1, 4	3
Ito et al. (92)	MC, RCT	Japan, 209	55.46	C/T	88.40	>18 yr, ANC < 1,000	FLCZ, 200 mg, p.o., QD	ITCZ, 200 mg, p.o., QD	1, 2, 4	3
Rijnders et al. (95)	DB, RCT	Netherlands, 271	49.49	C/T, 68.2%	48.70	ANC < 500, >10 days	aAMB, 12.5 mg, iNH, QD	Placebo	1, 3	7
Hiramatsu et al. (94)	OP, RCT	Japan, 100	46.90	SCT	19	≥18 yr, ANC ≤ 500/ μ l	FLCZ, 400 mg, i.v., QD	MCFG, 150 mg, i.v., QD	1, 2, 3, 4	6
Sawada et al. (96)	RCT	Japan, 107	6.01	C/T, 76.6%	27	<18 yr, ANC < 500	FLCZ, 10 mg/kg, i.v., QD	MCFG, 2 mg/kg, i.v., QD	1, 2, 4	4
Mattuzzi et al. (89)	OP, RCT	USA, 123	59.42	C/T	55.30	>18 yr, ANC < 500, >5 days	VOZC, 400 mg, i.v., QD	ITCZ, 200 mg, i.v., BID	1, 3, 4	5
Wingard et al. (28)	DB, MC, RCT	USA, 600	43.00	NA	76.50	>2 yr, ANC < 500, >5 days	FLCZ, 200 mg, p.o., BID	VOZC, 200 mg, p.o., BID	1, 2, 3	6

(Continued on next page)

TABLE 1 (Continued)

Reference	Trial design	Country, total no. of cases	Mean age (yr)	Main therapy	Leukemia (%)	Inclusion criteria ^c	Treatment arm ^e		Involved results ^d	Quality (Cochrane risk of bias)
							A	B		
Marks et al. (97)	OP, MC, RCT	International, 465	0.00	C/T	66.20	>12 yr, ANC < 500, >7 days	VOCZ, 200 mg, p.o., BID	ITCZ, 200 mg, p.o., BID	1, 2, 3, 4	5
Chaftari et al. (98)	OP, RCT	USA, 40	55.48	SCT	NA	>18 yr, ANC < 500, >7 days	p.o.CZ, 200 mg, p.o., TID	ABLC, 7.5 mg/kg, i.v., QD	1, 2, 4	3
Huang et al. (99)	OP, MC, RCT	China, 283	32.72	SCT	49	≥18 yr, ANC ≤ 500	ITCZ, 5 mg/kg, p.o., QD	MCFG, 50 mg, i.v., QD	1, 2, 4	5
Shen et al. (35)	OP, MC, RCT	China, 234	40.00	NA	88.00	>18 yr, ANC < 500, >7 days	p.o.CZ, 200 mg, p.o., TID	FLCZ, 400 mg, p.o., QD	1, 3, 4	5
Park et al. (100)	OP, RCT	South Korea, 250	46.66	SCT	40.80	>20 yr, ANC < 1,000, >5 days	FLCZ, 400 mg, p.o., QD	MFCG, 50 mg, i.v., QD	1, 2, 3, 4	3
Mahmoud et al. (19)	OP, RCT	Egypt, 70	7.35	C/T	NA	<18 yr, ANC < 500, >7 days	VOCZ, 4 mg/kg, i.v., BID	MCFG, 50 mg, i.v., QD	1, 2, 3	3

^aC/T, chemotherapy; SCT, stem cell transplant; MC, multi-center; DB, double blind; OP, open label; NA, not applicable.

^bCornely et al. (27) had three treatment arms: (i) p.o.CZ, 200 mg, p.o., TID; (ii) FLCZ, 400 mg, p.o., QD; (iii) ITCZ, 200 mg, p.o., BID.

^cInclusion criteria included age, ANC definition, and neutropenia duration.

^dInvolved results included the following groups: (1) invasive fungal infection, overall; (2) fungal infection, proven; (3) mortality, all cause; (4) adverse event, all causes.

^ep.o., per os; i.v., *in vitro*; QD, once a day; BID, twice a day; TID, three times a day.

^fPercentages indicate the percentages of patients receiving the indicated main therapy.




FIG 2 Schematic of the network of evidence used in network meta-analysis. Directly comparable treatments are linked with a line. iAMB, intravenous conventional amphotericin B; aAMB, aerosol amphotericin B; LAMB, liposomal amphotericin B; ABLC, amphotericin B lipid complex; ABCD, amphotericin B colloidal dispersion.

Furthermore, we conducted subgroup analyses to explore the potential spectrum of prophylaxis for invasive aspergillosis and invasive candidiasis. Posaconazole was also considered the superior choice to prevent invasive aspergillosis (RR, 0.13; 95% CI, 0.03 to 0.65), with significance of results (see the URL mentioned above, page 11). Compared with placebo, micafungin, fluconazole, posaconazole, voriconazole, and itraconazole had no difference in decreasing the incidence of invasive candidiasis (see the URL mentioned above, page 11).

Proven invasive fungal infection. The clinical judgement of a probable fungal infection would be uncertain and inconsistent among different physicians. Proven fungal infection has a more specific definition, which includes invasive mold infection that is demonstrated as fungal elements in tissues, evidence of associated tissue damage on biopsy specimens, and a proven invasive yeast infection based on isolation of yeast in the culture of a sample obtained by a sterile procedure from a normally sterile site, usually blood.

Based on the network meta-analysis results of the overall proven fungal infections (48 trials, 11,050 cases), posaconazole, micafungin, itraconazole, and fluconazole showed promising reduction of proven fungal infections, with RRs ranging from 0.14 to 0.54 (Fig. 4A). Among them, posaconazole was ranked highest in prevention of proven fungal infections (RR, 0.14; 95% CI, 0.05 to 0.38). In terms of the potential prophylaxis efficacy for different fungal spectra, posaconazole was the most effective for reduction of proven aspergillosis (RR, 0.13; 95% CI, 0.03 to 0.65) (Fig. 4B). Itraconazole, caspofungin, fluconazole, and intravenous conventional amphotericin B showed significantly decreased incidences of proven candidiasis; among them, itraconazole was ranked highest (RR, 0.21; 95% CI, 0.11 to 0.39) (Fig. 4C).

Figure 5 demonstrates the scatterplot of cumulative probabilities of being the most effective prophylactic antifungal agent for both invasive fungal infections and proven fungal infections. The scatterplot indicates that posaconazole is the most effective.

All-cause mortality and adverse events in the overall population. A total of 38 studies (8 arms, 8,447 cases) recorded the all-cause mortality outcome. The appendix at <https://goo.gl/6AAXgq>, page 12, demonstrates that there was no significant difference in the all-cause mortality rate between the 10 antifungal agents and placebo. Based on





FIG 3 NMA results (presented as risk ratio) for invasive fungal infection overall. (A) Forest plot of invasive fungal infections. (B) Multiple treatment comparisons for incidence of invasive fungal infection using consistency analysis based on network. P score was determined by SUCRA (surface under the cumulative ranking curve). CI, confidence interval; iAMB, intravenous conventional amphotericin B; aAMB, aerosol amphotericin B; LAMB, liposomal amphotericin B; ABLC, amphotericin B lipid complex; KTCZ, ketoconazole; FLCZ, fluconazole; ITCZ, itraconazole; VOCZ, voriconazole; POCZ, posaconazole; MCFG, micafungin; CASP, caspofungin.


cumulative ranking, liposomal amphotericin B tended to be the better choice for reduction of mortality (RR, 0.44; 95% CI, 0.14 to 1.39), but the difference was not significant (see the URL mentioned above, page 12).

A total of 28 studies (7,872 cases) documented the all-cause adverse events outcome (see the URL mentioned above, page 13). According to the results of the network meta-analysis, there was no significant difference in the incidence of overall adverse events between the antifungal agents and placebo. Based on the ranking probability scores, micafungin tended to have the lowest rate of overall all-cause adverse events (RR, 0.83; 95% CI, 0.53 to 1.31), but the difference was not significant (see the URL mentioned above, page 13). Based on Common Terminology Criteria for Adverse Events (CTCAE) v4.03 (29), we did subgroup analyses of the mild (grades 1 and 2) and severe (grade >2) adverse events groups. Micafungin, posaconazole, and fluconazole had significantly lower incidences of mild adverse events than oral polyenes (RRs ranged from 0.62 to 0.65) (see the URL mentioned above, page 13). For the severe adverse events, there were no significant differences among the antifungal agents (see the URL mentioned above, page 13). Based on the cumulative ranking, micafungin tended to be the safest prophylactic antifungal agent with the lowest incidence of adverse events, no matter what the severity (see the URL mentioned above, page 13). From the limited data, we analyzed the RR of common hepatic impairment and gastrointestinal upset in antifungal agent prevention. In terms of hepatic impairment,

A Proven Fungal Infection, overall



B Proven Aspergillosis Infection



C Proven Candidiasis Infection




FIG 4 NMA results (presented as risk ratio) for proven fungal infection. (A) Forest plot of proven fungal infection. (B) Forest plot of proven aspergillosis infection. (C) Forest plot of proven candidiasis infection. P score was determined by SUCRA (surface under the cumulative ranking curve). CI, confidence interval; iAMB, intravenous conventional amphotericin B; aAMB, aerosol amphotericin B; LAMB, liposomal amphotericin B; ABLC, amphotericin B lipid complex.




FIG 5 Scatter plot presenting the cumulative probability of prophylactic antifungal agent for overall invasive fungal infection and proven fungal infections. x axis, the probability of prevention for overall invasive fungal infections; y axis, the probability of prevention for proven fungal infections. Blue color means the probability scores of IFIs and PFIs are both above 50%, and red color means neither are.

voriconazole had a significantly increased risk (RR, 2.39; CI, 1.43 to 3.98); fluconazole, micafungin, itraconazole, and posaconazole did not show differences from the placebo (see the URL mentioned above, page 17). Furthermore, fluconazole, micafungin, itraconazole, posaconazole, and voriconazole had no statistically significant difference in the incidence of gastrointestinal upset compared to that of the placebo (see the URL mentioned above, page 17).

Sensitivity test. Due to including extensive antifungal agent studies, we assessed whether the prophylactic effect for the primary outcome was robust in subgroup analyses and sensitivity test using study year, study quality, study precision (i.e., small study effect), and homophilic pattern in antifungal agent trials (30). The appendix summarizes the definitions of covariates, and the results did not change substantially (see the URL mentioned above, page 16). The preference for fluconazole comparisons and avoidance of head-to-head new drug comparisons in antifungal trials was pointed out by Rizos et al. (30). In our present research, 25 trials used comparisons with fluconazole, accounting for nearly 46% of the included studies. After excluding the studies of fluconazole comparisons, the sensitivity test showed no substantial difference for the original network meta-analysis. Although the values of relative risk were changed, posaconazole was ranked as the most effective, and micafungin, voriconazole, and caspofungin had similar cumulative ranking results (see the URL mentioned above, page 16). We used comparison-adjusted funnel plots to investigate whether results in imprecise trials differ from those in more precise trials. Net-funnel plot (see the URL mentioned above, page 14) and egger graph (see the URL mentioned above, page 14) demonstrated no significant publication bias. Furthermore, the design-by-treatment interaction model and side-splitting model reported no significant inconsistency.

DISCUSSION

There is an emerging need for synthesis of evidence on primary fungal prophylaxis in hematological malignancy patients with neutropenic fever, especially because a myriad of novel treatments has become available in the past decade. Several systematic reviews focused on separate antifungal agent comparisons. Only one network meta-analysis has synthesized the evidence on IFI prophylaxis in hematological cancer on the

basis of pooled odds ratios from randomized controlled trials (31). However, the evidence shown by Leonart et al. (31) was focused on double-blind trials; among 25 trials, there were 19 trials that compared a single antifungal with placebo instead of head-to-head studies. Currently, conclusions on the comparisons across prior network meta-analyses cannot be clearly made.

Our purpose was to pool all qualifying evidence and enable an integrated comparison of all current antifungal agents for IFI prevention in hematological malignancy patients with febrile neutropenia. In this study, RR was adopted as the effect measure for IFI incidence, more recent treatments were included, and all pieces of evidence were combined into a single network. Indeed, we were able to combine within one network the evidence from 54 enrolled RCTs on a total of 12,832 cases, including 13 treatment arms, using random-effect network meta-analysis. Our analyses provided crucial information on health care decision-making for primary fungal prophylaxis in such patients. Of the 13 treatment options, posaconazole was the top priority in our network meta-analysis, both in terms of ranking and probability of being the most effective treatment. Based on pooled evidence, posaconazole was ranked highest in reduction of the overall IFIs, including invasive aspergillosis. These results strongly supported the current Infectious Disease Society of America guidelines for primary prophylaxis in patients with acute leukemia and post-stem cell transplant (32). For proven fungal infections, the overall incidence was significantly reduced with posaconazole, micafungin, itraconazole, and fluconazole. Moreover, posaconazole was the better choice for prevention of proven aspergillosis. For proven candidiasis prophylaxis, itraconazole, fluconazole, caspofungin, and intravenous conventional amphotericin B presented with significantly good efficacy. For patients who live in regions with a high prevalence of candidiasis, the above-mentioned drugs would be the optimal choice.

We proved that posaconazole prophylaxis was associated with significant reductions in overall invasive, proven fungal infection and invasive aspergillosis but did not have a significant impact on all-cause mortality. These discordant results had been disclosed in clinical trials (33–35), meta-analyses (15, 16, 36, 37), and real-world data (38–40). The absence of a significant difference in all-cause mortality might be explained by the following hypotheses. The first possible explanation for the insignificant effect on all-cause mortality is that empirical treatment with voriconazole is associated with substantially improved prognosis of invasive aspergillosis (41, 42). The second possible reason for the discrepancy between significant reduction in the incidence of invasive fungal infection and the lack of impact on all-cause mortality is associated with the use of galactomannan tests. Marr et al. (43) reported that mold-active azole agents could interfere with the sensitivity of this test, and Kim et al. (44) showed micafungin would limit galactomannan detection, thus the true reduction of invasive fungal diseases observed with mold-active agents may be overestimated. The third possibility is that mold-active prophylaxis increases non-IFI-related deaths. It is also possible that drug interactions further contributed to increased patient deaths.

Our analysis on overall adverse events did not show any significant differences, except for the use of conventional iAMB, which was significantly ranked at the bottom of the list. Subgroup analyses revealed that micafungin had a significantly lower rate of mild adverse events, which may be a valuable clinical concern for rotating antifungal agents. These results were somewhat in line with clinical experience and earlier observations (45).

All available prophylaxis options in hematological cancer were systematically obtained from the published domain and were pooled in a conventional and mathematically transparent way. Therefore, this systematic literature review and network meta-analysis are reproducible and could compare all treatment options that are currently available for primary prophylaxis in such patients.

To examine the potential biases in antifungal agent studies, we assessed whether the prophylactic effect for the primary outcome was robust in subgroup analyses and sensitivity tests. The important variables of quality scores, sample sizes, publication years, and preference of fluconazole pattern were detected. No substantial biases existed in the sensitivity test. Consequently, with included emerging studies comparing

new drugs to each other, our present network meta-analysis is considered more robust. However, we did not consider differences in dosing schemes and modes of administration; among the studies, the most apparent were changes in the mode of fluconazole and itraconazole administration from oral to intravenous due to intolerance. However, in clinical practice, shifting from oral to intravenous administration and differences in bioavailability are probably common. Therefore, differences in dosing schemes and modes of administration remain a subject of further research. Readers should be aware that our results, similar to the results of each trial, apply to the average patient. Although this study might assist doctors and patients in evidence-based decision-making, selecting the most appropriate treatment option also depends on the individual patient characteristics and preferences.

With our best effort, we provided evidence that in hematological malignancy patients who develop neutropenic fever after myelosuppressive chemotherapy or hematopoietic cell transplantation, posaconazole is a somewhat superior option for primary fungal prophylaxis. For patients living in areas with a high incidence of candidiasis infection, micafungin, itraconazole, and fluconazole would be more effective options. When there is a need to rotate antifungal agents due to adverse events, micafungin would be a safer choice. Despite the limitations of this study, our results provided insight on the rank order of efficacy and safety of these treatment options. This can facilitate evidence-based decision-making in the clinical setting, where most treatment agents have not been compared in head-to-head RCT settings or have not been previously compared using evidence synthesis. Nevertheless, we emphasize that it remains essential to conduct phase III trials to obtain more direct head-to-head evidence. Until such evidence becomes available, our results are highly important for informed decision-making in everyday clinical practice.

MATERIALS AND METHODS

We conducted a network meta-analysis in a frequentist framework and used the *Cochrane Handbook for Systematic Reviews of Interventions* approach to evaluate the quality of evidence (46). This study has been registered at PROSPERO (<http://www.crd.york.ac.uk/PROSPERO>) with registration number CRD42017058429.

Search methods and criteria for considering studies for inclusion. We performed an extensive electronic search (see the appendix at <https://goo.gl/6AAXGq>, pages 2 to 6) of the Cochrane Central Register of Controlled Trials, PubMed, and Embase for RCTs. We further sought details of the trials or protocols from <https://clinicaltrials.gov> to establish the eligibility of potential trials. No language restrictions were applied. Our latest search was completed on 19 February 2017. Two reviewers (C.H.L. and C.L.) screened the titles and abstracts of the retrieved articles to search for the eligible RCTs.

Types of studies. All available RCTs that compared the prophylactic efficacy and safety of antifungal agents in hematological cancer were enrolled. Placebo-designed studies were included to serve as a reference comparator for estimating the relative effectiveness of antifungal agents. In the present study, we assumed no relative differences in effectiveness between placebo and no prophylaxis. RCTs with quasi-experiment and crossover designs were excluded. Study design of comparison of different dosage, the comparator of historical control, intervention with non-FDA-approved antifungal agents, and unclear control groups (like standard policy) were also excluded.

Types of participants. We enrolled patients, of any age, with a diagnosis of hematological malignancy and who were under myelosuppressive chemotherapy or hematopoietic cell transplantation and developed neutropenic fever, which was defined as an absolute neutrophil count of $<1,500/\mu\text{l}$ for at least 3 days. Autogenic or allogeneic stem cell transplants were not limited. Those who received concomitant antibacterial agents were included. Patients who received antifungal agents for prophylaxis or empirical treatment prior to enrollment in the trial were excluded.

Types of interventions and comparison. The trials included were those that evaluated the prophylactic efficacy and safety of polyenes, azoles, and echinocandins. The traditional amphotericin B formulations are now prescribed less frequently due to drug toxicity, and these were designated the reference comparator because of several studies of head-to-head comparisons with other antifungal agents in past years. Furthermore, we also chose to enroll trials involving oral polyene and ketoconazole because, despite not being recommended in clinical practice any longer for invasive fungal infection prophylaxis, we did not want to lose any evidence from the literature. The dosage and duration of the prophylactic or antifungal agent were not limited.

Types of outcomes. RCTs that assessed the efficacy of primary fungal prophylaxis using the following outcomes were included: (i) IFI incidence rates (the classification criteria for possible, probable, and proven fungal infections were based on the 2008 criteria by the EORTC and MSG [47]), (ii) all-cause or drug-related mortality, and (iii) all-cause or drug-related adverse events (defined based on the Common Terminology Criteria of Adverse Events, v4.03) (29). We chose the longest follow-up time (end of follow-up or death) as the measurement time point for all outcomes.

Risk of bias assessment. The study quality was assessed by two reviewers (C.H.L. and C.L.) using the methodology and categories described in the Cochrane Collaboration handbook (46). In case of disagreement, a group discussion was done to reach a consensus. In the assessment of other issues, we focused on the baseline imbalance and source of financial support (48).

Data extraction. Two reviewers (C.H.L. and C.L.) independently assessed the eligibility of all identified citations and extracted data from original trial reports using a specifically designed form that captured information on the study characteristics, patient characteristics, and sample sizes and details of interventions with comparisons and outcomes. To lower the chances of entry error, double data entry and cross-checking were performed.

Data synthesis and statistical analysis. A network meta-analysis was conducted to simultaneously compare 13 antifungal treatment options for each outcome. The network meta-analysis in the present study was performed by using the netmeta package (0.9-7) in R, version 3.4.1 (49). The incidence of IFIs was defined as the primary outcome. RR with 95% CI was calculated using the random-effect model by following UK NICE guidelines (50). A *P* value of less than 0.05 was considered statistically significant.

A network plot that represented the overall information of the trials included in the analysis was generated (51). The contribution of each direct comparison to each network estimate was calculated according to the variance of the direct treatment effect and the network structure (52). Network meta-analysis is a method for global assessment of the current evidence; therefore, we needed to carefully compare indirect evidence with the direct estimates. Inconsistency referred to the differences between the various direct and indirect effects that were estimated for the same comparison. Inconsistency was evaluated by design-by-treatment interaction model and side-splitting model.

A forest plot of the estimated summary effects, along with the CI for all comparisons, was generated to summarize the relative mean effects, the predictions on each comparison in one plot (53). We estimated the probability of a treatment being ranked at a specific place according to the outcome (*P* score) using surface under the cumulative ranking curve (SUCRA), which is a simple transformation of the mean rank to provide a hierarchy of the treatments and to account for the location and variance of all relative treatment effects. A higher SUCRA value reflected a higher possible ranking of the treatment (54). The *P* score represented the SUCRA value in the forest plot.

REFERENCES

- Hamza NS, Ghannoum MA, Lazarus HM. 2004. Choices aplenty: antifungal prophylaxis in hematopoietic stem cell transplant recipients. *Bone Marrow Transplant* 34:377–389. <https://doi.org/10.1038/sj.bmt.1704603>.
- Baddley JW, Andes DR, Marr KA, Kontoyiannis DP, Alexander BD, Kauffman CA, Oster RA, Anaissie EJ, Walsh TJ, Schuster MG, Wingard JR, Patterson TF, Ito JI, Williams OD, Chiller T, Pappas PG. 2010. Factors associated with mortality in transplant patients with invasive aspergillosis. *Clin Infect Dis* 50:1559–1567. <https://doi.org/10.1086/652768>.
- Goodman JL, Winston DJ, Greenfield RA, Chandrasekar PH, Fox B, Kaizer H, Shadduck RK, Shea TC, Stiff P, Friedman DJ, Powderly WG, Silber JL, Horowitz H, Lichtin L, Wolff SN, Mangan KF, Silver SM, Daniel Weisdorf D, Ho WG, Gilbert G, Buell D. 1992. A controlled trial of fluconazole to prevent fungal infections in patients undergoing bone marrow transplantation. *N Engl J Med* 326:845–851. <https://doi.org/10.1056/NEJM199203263261301>.
- Winston DJ, Maziarz RT, Chandrasekar PH, Lazarus HM, Goldman M, Blumer JL, Leitz GJ, Territo MC. 2003. Intravenous and oral itraconazole versus intravenous and oral fluconazole for long-term antifungal prophylaxis in allogeneic hematopoietic stem-cell transplant recipients. A multicenter, randomized trial. *Ann Intern Med* 138:705–713. <https://doi.org/10.7326/0003-4819-138-9-200305060-00006>.
- El-Cheikh J, Faucher C, Furst S, Duran S, Berger P, Vey N, Stoppa AM, Bouabdallah R, Gastaut JA, Viens P, Blaise D, Mohty M. 2007. High-dose weekly liposomal amphotericin B antifungal prophylaxis following reduced-intensity conditioning allogeneic stem cell transplantation. *Bone Marrow Transplant* 39:301–306. <https://doi.org/10.1038/sj.bmt.1705592>.
- Rogers TR, Slavin MA, Donnelly JP. 2011. Antifungal prophylaxis during treatment for hematological malignancies: are we there yet? *Br J Haematol* 153:681–697. <https://doi.org/10.1111/j.1365-2141.2011.08650.x>.
- Leventakos K, Lewis RE, Kontoyiannis DP. 2010. Fungal infections in leukemia patients: how do we prevent and treat them? *Clin Infect Dis* 50:405–415. <https://doi.org/10.1086/649879>.
- Tacke D, Buchheidt D, Karthaus M, Krause SW, Maschmeyer G, Neumann S, Ostermann H, Penack O, Rieger C, Ruhnke M, Sandherr M, Schweer KE, Ullmann AJ, Cornely OA. 2014. Primary prophylaxis of invasive fungal infections in patients with hematologic malignancies. 2014 update of the recommendations of the Infectious Diseases Working Party of the German Society for Hematology and Oncology. *Ann Hematol* 93:1449–1456. <https://doi.org/10.1007/s00277-014-2108-y>.
- Powles R, Mehta J, Kulkarni S, Treleaven J, Millar B, Marsden J, Shepherd V, Rowland A, Sirohi B, Tait D, Horton C, Long S, Singhal S. 2000. Allogeneic blood and bone-marrow stem-cell transplantation in hematological malignant diseases: a randomised trial. *Lancet* 355:1231–1237. [https://doi.org/10.1016/S0140-6736\(00\)02090-0](https://doi.org/10.1016/S0140-6736(00)02090-0).
- Marr KA, Carter RA, Boeckh M, Martin P, Corey L. 2002. Invasive aspergillosis in allogeneic stem cell transplant recipients: changes in epidemiology and risk factors. *Blood* 100:4358–4366. <https://doi.org/10.1182/blood-2002-05-1496>.
- Neofytos D, Horn D, Anaissie E, Steinbach W, Olyaei A, Fishman J, Pfaller M, Chang C, Webster K, Marr K. 2009. Epidemiology and outcome of invasive fungal infection in adult hematopoietic stem cell transplant recipients: analysis of Multicenter Prospective Antifungal Therapy (PATH) Alliance registry. *Clin Infect Dis* 48:265–273. <https://doi.org/10.1086/595846>.
- Kontoyiannis DP, Marr KA, Park BJ, Alexander BD, Anaissie EJ, Walsh TJ, Ito J, Andes DR, Baddley JW, Brown JM, Brumble LM, Freifeld AG, Hadley S, Herwaldt LA, Kauffman CA, Knapp K, Lyon GM, Morrison VA, Papanicolaou G, Patterson TF, Perl TM, Schuster MG, Walker R, Wanne-muehler KA, Wingard JR, Chiller TM, Pappas PG. 2010. Prospective surveillance for invasive fungal infections in hematopoietic stem cell transplant recipients, 2001–2006: overview of the Transplant-Associated Infection Surveillance Network (TRANSNET) Database. *Clin Infect Dis* 50:1091–1100. <https://doi.org/10.1086/651263>.
- Even C, Bastuji-Garin S, Hicheri Y, Pautas C, Botterel F, Maury S, Cabanne L, Bretagne S, Cordonnier C. 2011. Impact of invasive fungal disease on the chemotherapy schedule and event-free survival in acute leukemia patients who survived fungal disease: a case-control study. *Haematologica* 96:337–341. <https://doi.org/10.3324/haematol.2010.030825>.
- Lewis RE, Cahyame-Zuniga L, Leventakos K, Chamilos G, Ben-Ami R, Tamboli P, Tarrand J, Bodey GP, Luna M, Kontoyiannis DP. 2013. Epidemiology and sites of involvement of invasive fungal infections in patients with hematological malignancies: a 20-year autopsy study. *Mycoses* 56:638–645. <https://doi.org/10.1111/myc.12081>.
- Robenshtok E, Gafter-Gvili A, Goldberg E, Weinberger M, Yeshurun M, Leibovici L, Paul M. 2007. Antifungal prophylaxis in cancer patients after chemotherapy or hematopoietic stem-cell transplantation: systematic review and meta-analysis. *J Clin Oncol* 25:5471–5489. <https://doi.org/10.1200/JCO.2007.12.3851>.
- Ethier MC, Science M, Beyene J, Briel M, Lehmbecher T, Sung L. 2012.

- Mould-active compared with fluconazole prophylaxis to prevent invasive fungal diseases in cancer patients receiving chemotherapy or haematopoietic stem-cell transplantation: a systematic review and meta-analysis of randomised controlled trials. *Br J Cancer* 106:1626–1637. <https://doi.org/10.1038/bjc.2012.147>.
17. Uzun O, Anaissie EJ. 1995. Antifungal prophylaxis in patients with hematologic malignancies: a reappraisal. *Blood* 86:2063–2072.
 18. Jeong SH, Kim DY, Jang JH, Mun YC, Choi CW, Kim SH, Kim JS, Park JS. 2016. Efficacy and safety of micafungin versus intravenous itraconazole as empirical antifungal therapy for febrile neutropenic patients with hematological malignancies: a randomized, controlled, prospective, multicenter study. *Ann Hematol* 95(2):337–344. <https://doi.org/10.1007/s00277-015-2545-2>.
 19. Mahmoud AM, Shalaby LM, El-Malhallowy HA, AhmadKhalil S. 2016. Micafungin as a prophylactic antifungal during induction phase of chemotherapy for pediatric acute myeloblastic leukemia. MSc degree in Pediatric Oncology, Cairo University, Cairo, Egypt. <https://goo.gl/oqSVsh>.
 20. Oyake T, Kowata S, Murai K, Ito S, Akagi T, Kubo K, Sawada K, Ishida Y. 2016. Comparison of micafungin and voriconazole as empirical antifungal therapies in febrile neutropenic patients with hematological disorders: a randomized controlled trial. *Eur J Haematol* 96(6):602–609. <https://doi.org/10.1111/ejh.12641>.
 21. Park S, Kim K, Jang JH, Kim SJ, Kim WS, Chung DR, Kang CI, Peck KR, Jung CW. 2016. Randomized trial of micafungin versus fluconazole as prophylaxis against invasive fungal infections in hematopoietic stem cell transplant recipients. *J Infect* 73:496–505. <https://doi.org/10.1016/j.jinf.2016.06.011>.
 22. Bow EJ, Vannest DJ, Slavin M, Cordonnier C, Cornely OA, Marks DI, Pagliuca A, Solano C, Cragin L, Shaul AJ, Sorensen S, Chambers R, Kantecki M, Weinstein D, Schlamm H. 2015. Systematic review and mixed treatment comparison meta-analysis of randomized clinical trials of primary oral antifungal prophylaxis in allogeneic hematopoietic cell transplant recipients. *BMC Infect Dis* 15:128. <https://doi.org/10.1186/s12879-015-0855-6>.
 23. Wang JF, Xue Y, Zhu XB, Fan H. 2015. Efficacy and safety of echinocandins versus triazoles for the prophylaxis and treatment of fungal infections: a meta-analysis of RCTs. *Eur J Clin Microbiol Infect Dis* 34:651–659. <https://doi.org/10.1007/s10096-014-2287-4>.
 24. Xu SX, Shen JL, Tang XF, Feng B, Xu HQ. 2016. Newer antifungal agents micafungin and voriconazole for fungal infection prevention during hematopoietic cell transplantation: a meta-analysis. *Eur Rev Med Pharmacol Sci* 20:381–390.
 25. Lu G, Ades AE. 2004. Combination of direct and indirect evidence in mixed treatment comparisons. *Stat Med* 23:3105–3124. <https://doi.org/10.1002/sim.1875>.
 26. Jansen JP, Crawford B, Bergman G, Stam W. 2008. Bayesian meta-analysis of multiple treatment comparisons: an introduction to mixed treatment comparisons. *Value Health* 11:956–964. <https://doi.org/10.1111/j.1524-4733.2008.00347.x>.
 27. Cornely OA, Maertens J, Winston DJ, Perfect J, Ullmann AJ, Walsh TJ, Helfgott D, Holowiecki J, Stockelberg D, Goh YT, Petrini M, Hardalo C, Suresh R, Angulo-Gonzalez D. 2007. Posaconazole vs. fluconazole or itraconazole prophylaxis in patients with neutropenia. *N Engl J Med* 356:348–359. <https://doi.org/10.1056/NEJMoa061094>.
 28. Wingard JR, Carter SL, Walsh TJ, Kurtzberg J, Small TN, Baden LR, Gersten ID, Mendizabal AM, Leather HL, Confer DL, Maziarz RT, Stadtmauer EA, Bolanos-Meade J, Brown J, Dipersio JF, Boeckh M, Marr KA, Blood Marrow Transplant Clinical Trials Network. 2010. Randomized, double-blind trial of fluconazole versus voriconazole for prevention of invasive fungal infection after allogeneic hematopoietic cell transplantation. *Blood* 116:5111–5118. <https://doi.org/10.1182/blood-2010-02-268151>.
 29. Zhang S, Liang F, Tannock I. 2016. Use and misuse of common terminology criteria for adverse events in cancer clinical trials. *BMC Cancer* 16:392. <https://doi.org/10.1186/s12885-016-2408-9>.
 30. Rizos EC, Salanti G, Kontoyiannis DP, Ioannidis JP. 2011. Homophily and co-occurrence patterns shape randomized trials agendas: illustration in antifungal agents. *J Clin Epidemiol* 64:830–842. <https://doi.org/10.1016/j.jclinepi.2010.11.017>.
 31. Leonart LP, Tonin FS, Ferreiro VL, Penteado STS, Wiens A, Motta FA, Pontarolo R. 2017. A network meta-analysis of primary prophylaxis for invasive fungal infection in hematological patients. *J Clin Pharm Ther* 42:530–538. <https://doi.org/10.1111/jcpt.12579>.
 32. Patterson TF, Thompson GR, III, Denning DW, Fishman JA, Hadley S, Herbrecht R, Kontoyiannis DP, Marr KA, Morrison VA, Nguyen MH, Segal BH, Steinbach WJ, Stevens DA, Walsh TJ, Wingard JR, Young JA, Bennett JE. 2016. Executive summary: practice guidelines for the diagnosis and management of aspergillosis: 2016 update by the Infectious Diseases Society of America. *Clin Infect Dis* 63:433–442. <https://doi.org/10.1093/cid/ciw444>.
 33. Winston DJ, Chandrasekar PH, Lazarus HM, Goodman JL, Silber JL, Horowitz H, Shadduck RK, Rosenfeld CS, Ho WG, Islam MZ, Buell DN. 1993. Fluconazole prophylaxis of fungal infections in patients with acute leukemia: results of a randomized placebo-controlled, double-blind, multicenter trial. *Ann Intern Med* 118:495–503. <https://doi.org/10.7326/0003-4819-118-7-199304010-00003>.
 34. Ullmann AJ, Lipton JH, Vesole DH, Chandrasekar P, Langston A, Taranoto SR, Greinix H, Morais de Azevedo W, Reddy V, Boparai N, Pedicone L, Patino H, Durrant S. 2007. Posaconazole or fluconazole for prophylaxis in severe graft-versus-host disease. *N Engl J Med* 356:335–347. <https://doi.org/10.1056/NEJMoa061098>.
 35. Shen Y, Huang XJ, Wang JX, Jin J, Hu JD, Yu K, Wu DP, Wang SJ, Yu L, Chen XQ, Liu T, Liang YM, Chen FP, Li Y, Shen ZX. 2013. Posaconazole vs. fluconazole as invasive fungal infection prophylaxis in China: a multicenter, randomized, open-label study. *Int J Clin Pharmacol Ther* 51:738–745. <https://doi.org/10.5414/CP201880>.
 36. Vardakas KZ, Michalopoulos A, Falagas ME. 2005. Fluconazole versus itraconazole for antifungal prophylaxis in neutropenic patients with hematological malignancies: a meta-analysis of randomized-controlled trials. *Br J Haematol* 131:22–28. <https://doi.org/10.1111/j.1365-2141.2005.05727.x>.
 37. Wang J, Zhan P, Zhou R, Xu J, Shao X, Yang Y, Ouyang J. 2010. Prophylaxis with itraconazole is more effective than prophylaxis with fluconazole in neutropenic patients with hematological malignancies: a meta-analysis of randomized-controlled trials. *Med Oncol* 27: 1082–1088. <https://doi.org/10.1007/s12032-009-9339-0>.
 38. Ananda-Rajah MR, Grigg A, Downey MT, Bajel A, Spelman T, Cheng A, Thursky KT, Vincent J, Slavin MA. 2012. Comparative clinical effectiveness of prophylactic voriconazole/posaconazole to fluconazole/itraconazole in patients with acute myeloid leukemia/myelodysplastic syndrome undergoing cytotoxic chemotherapy over a 12-year period. *Haematologica* 97:459–463. <https://doi.org/10.3324/haematol.2011.051995>.
 39. Kung HC, Johnson MD, Drew RH, Saha-Chaudhuri P, Perfect JR. 2014. Clinical effectiveness of posaconazole versus fluconazole as antifungal prophylaxis in hematology-oncology patients: a retrospective cohort study. *Cancer Med* 3:667–673. <https://doi.org/10.1002/cam4.225>.
 40. Dahlen T, Kalin M, Cederlund K, Nordlander A, Bjorkholm M, Ljungman P, Blennow O. 2016. Decreased invasive fungal disease but no impact on overall survival by posaconazole compared to fluconazole prophylaxis: a retrospective cohort study in patients receiving induction therapy for acute myeloid leukaemia/myelodysplastic syndromes. *Eur J Haematol* 96:175–180. <https://doi.org/10.1111/ejh.12565>.
 41. Herbrecht R, Denning DW, Patterson TF, Bennett JE, Greene RE, Oestmann JW, Kern WV, Marr KA, Ribaud P, Lortholary O, Sylvester R, Rubin RH, Wingard JR, Stark P, Durand C, Caillot D, Thiel E, Chandrasekar PH, Hodges MR, Schlamm HT, Troke PF, de Pauw B, Invasive Fungal Infections Group of the European Organisation for Research and Treatment of Cancer and the Global Aspergillus Study Group. 2002. Voriconazole versus amphotericin B for primary therapy of invasive aspergillosis. *N Engl J Med* 347:408–415. <https://doi.org/10.1056/NEJMoa020191>.
 42. Nivoix Y, Velten M, Letscher-Bru V, Moghaddam A, Natarajan-Ame S, Fohrer C, Lioure B, Bilger K, Lutun P, Marcellin L, Launoy A, Freys G, Bergerat JP, Herbrecht R. 2008. Factors associated with overall and attributable mortality in invasive aspergillosis. *Clin Infect Dis* 47: 1176–1184. <https://doi.org/10.1086/592255>.
 43. Marr KA, Laverdiere M, Gugel A, Leisenring W. 2005. Antifungal therapy decreases sensitivity of the Aspergillus galactomannan enzyme immunoassay. *Clin Infect Dis* 40:1762–1769. <https://doi.org/10.1086/429921>.
 44. Kim R, Koh Y, Shin DY, Choe PG, Kim NJ, Yoon SS, Oh MD, Park WB, Kim I. 2017. The limited role of serum galactomannan assay in screening for invasive pulmonary aspergillosis in allogeneic stem cell transplantation recipients on micafungin prophylaxis: a retrospective study. *Blood Res* 52:300–306. <https://doi.org/10.5045/br.2017.52.4.300>.
 45. Lee CH, Lin JC, Ho CL, Sun M, Yen WT, Lin C. 2017. Efficacy and safety of micafungin versus extensive azoles in the prevention and treatment of invasive fungal infections for neutropenia patients with hematolog-

- ical malignancies: a meta-analysis of randomized controlled trials. *PLoS One* 12:e0180050. <https://doi.org/10.1371/journal.pone.0180050>.
46. Higgins JPT, Green S (ed). 2013. *Cochrane handbook for systematic reviews of interventions* version 5.1.0. The Cochrane Collaboration. <http://handbook.cochrane.org/>.
 47. De Pauw B, Walsh TJ, Donnelly JP, Stevens DA, Edwards JE, Calandra T, Pappas PG, Maertens J, Lortholary O, Kauffman CA, Denning DW, Patterson TF, Maschmeyer G, Bille J, Dismukes WE, Herbrecht R, Hope WW, Kibbler CC, Kullberg BJ, Marr KA, Munoz P, Odds FC, Perfect JR, Restrepo A, Ruhnke M, Segal BH, Sobel JD, Sorrell TC, Viscoli C, Wingard JR, Zaoutis T, Bennett JE. 2008. Revised definitions of invasive fungal disease from the European Organization for Research and Treatment of Cancer/Invasive Fungal Infections Cooperative Group and the National Institute of Allergy and Infectious Diseases Mycoses Study Group (EORTC/MSG) Consensus Group. *Clin Infect Dis* 46:1813–1821. <https://doi.org/10.1086/588660>.
 48. Roberts C, Torgerson DJ. 1999. Understanding controlled trials: baseline imbalance in randomised controlled trials. *BMJ* 319:185.
 49. Rücker GS, Krahn U, König J. 2017. Package netmeta: network meta-analysis using frequentist methods (R package version 0.9-7.). <https://CRAN.R-project.org/package=netmeta>. Accessed 4 August 2017.
 50. Leyva F, Plummer CJ. 2015. National Institute for Health and Care Excellence 2014 guidance on cardiac implantable electronic devices: health economics reloaded. *Europace* 17:339–342. <https://doi.org/10.1093/europace/euu384>.
 51. Salanti G. 2012. Indirect and mixed-treatment comparison, network, or multiple-treatments meta-analysis: many names, many benefits, many concerns for the next generation evidence synthesis tool. *Res Synth Methods* 3:80–97. <https://doi.org/10.1002/jrsm.1037>.
 52. Krahn U, Binder H, König J. 2013. A graphical tool for locating inconsistency in network meta-analyses. *BMC Med Res Methodol* 13:35. <https://doi.org/10.1186/1471-2288-13-35>.
 53. Jackson D, White IR, Riley RD. 2012. Quantifying the impact of between-study heterogeneity in multivariate meta-analyses. *Stat Med* 31:3805–3820. <https://doi.org/10.1002/sim.5453>.
 54. Riley RD, Higgins JP, Deeks JJ. 2011. Interpretation of random effects meta-analyses. *BMJ* 342:d549. <https://doi.org/10.1136/bmj.d549>.
 55. Donnelly JP, Starke ID, Galton DA, Catovsky D, Goldman JM, Darrell JH. 1984. Oral ketoconazole and amphotericin B for the prevention of yeast colonization in patients with acute leukaemia. *J Hosp Infect* 5:83–91. [https://doi.org/10.1016/0195-6701\(84\)90105-1](https://doi.org/10.1016/0195-6701(84)90105-1).
 56. Palmblad J, Lonnqvist B, Carlsson B, Grimfors G, Jarnmark M, Lerner R, Ljungman P, Nystrom-Rosander C, Petrini B, Oberg G. 1992. Oral ketoconazole prophylaxis for *Candida* infections during induction therapy for acute leukaemia in adults: more bacteraemias. *J Intern Med* 231:363–370. <https://doi.org/10.1111/j.1365-2796.1992.tb00945.x>.
 57. Akiyama H, Mori S, Tanikawa S, Sakamaki H, Onozawa Y. 1993. Fluconazole versus oral amphotericin B in preventing fungal infection in chemotherapy-induced neutropenic patients with haematological malignancies. *Mycoses* 36:373–378. <https://doi.org/10.1111/j.1439-0507.1993.tb00725.x>.
 58. Philpott-Howard JN, Wade JJ, Mufti GJ, Brammer KW, Ehninger G. 1993. Randomized comparison of oral fluconazole versus oral polyenes for the prevention of fungal infection in patients at risk of neutropenia. Multicentre Study Group. *J Antimicrob Chemother* 31:973–984. <https://doi.org/10.1093/jac/31.6.973>.
 59. Vreugdenhil G, Van Dijke BJ, Donnelly JP, Novakova IRO, Raemaekers JMM, Hoogkamp-Korstanje MAA, Koster M, De Pauw BE. 1993. Efficacy of itraconazole in the prevention of fungal infections among neutropenic patients with hematologic malignancies and intensive chemotherapy: a double blind, placebo controlled study. *Leuk Lymphoma* 11:353–358. <https://doi.org/10.3109/10428199309067926>.
 60. Bodey GP, Anaissie EJ, Elting LS, Estey E, O'Brien S, Kantarjian H. 1994. Antifungal prophylaxis during remission induction therapy for acute leukemia fluconazole versus intravenous amphotericin B. *Cancer* 73:2099–2106. [https://doi.org/10.1002/1097-0142\(19940415\)73:8<2099::AID-CNCR2820730814>3.0.CO;2-N](https://doi.org/10.1002/1097-0142(19940415)73:8<2099::AID-CNCR2820730814>3.0.CO;2-N).
 61. Chandrasekar PH, Gatny CM. 1994. The effect of fluconazole prophylaxis on fungal colonization in neutropenic cancer patients. *J Antimicrob Chemother* 33:309–318. <https://doi.org/10.1093/jac/33.2.309>.
 62. Ellis ME, Clink H, Ernst P, Halim MA, Padmos A, Spence D, Kalin M, Hussein Qadri SM, Burnie J, Greer W. 1994. Controlled study of fluconazole in the prevention of fungal infections in neutropenic patients with haematological malignancies and bone marrow transplant recipients. *Eur J Clin Microbiol Infect Dis* 13:3–11. <https://doi.org/10.1007/BF02026116>.
 63. Ninane J. 1994. A multicentre study of fluconazole versus oral polyenes in the prevention of fungal infection in children with hematological or oncological malignancies. Multicentre Study Group. *Eur J Clin Microbiol Infect Dis* 13:330–337. <https://doi.org/10.1007/BF01974614>.
 64. Riley DK, Pavia AT, Beatty PG, Petersen FB, Spruance JL, Stokes R, Evans TG. 1994. The prophylactic use of low-dose amphotericin B in bone marrow transplant patients. *Am J Med* 97:509–514. [https://doi.org/10.1016/0002-9343\(94\)90345-X](https://doi.org/10.1016/0002-9343(94)90345-X).
 65. Annaloro C, Oriana A, Tagliaferri E, Bertolli V, Della Volpe A, Soligo D, Ibatici A, Pozzoli E, Lambertenghi Deliliers GL. 1995. Efficacy of different prophylactic antifungal regimens in bone marrow transplantation. *Haematologica* 80:512–517.
 66. Behre GF, Schwartz S, Lenz K, Ludwig WD, Wandt H, Schilling E, Heinemann V, Link H, Trittin A, Boenisch O, Treder W, Siebert W, Hiddemann W, Beyer J. 1995. Aerosol amphotericin B inhalations for prevention of invasive pulmonary aspergillosis in neutropenic cancer patients. *Ann Hematol* 71:287–291. <https://doi.org/10.1007/BF01697981>.
 67. Egger T, Gratwohl A, Tichelli A, Uhr M, Stebler Gysi C, Passweg J, Pless M, Wernli M, Buser U, Wuhmann J. 1995. Comparison of fluconazole with oral polyenes in the prevention of fungal infections in neutropenic patients. A prospective, randomized, single-center study. *Support Care Cancer* 3:139–146. <https://doi.org/10.1007/BF00365855>.
 68. Schaffner A, Schaffner M. 1995. Effect of prophylactic fluconazole on the frequency of fungal infections, amphotericin B use, and health care costs in patients undergoing intensive chemotherapy for hematologic neoplasias. *J Infect Dis* 172:1035–1041. <https://doi.org/10.1093/infdis/172.4.1035>.
 69. Kern W, Behre G, Rudolf T, Kerkhoff A, Grote-Metke A, Eimermacher H, Kubica U, Wormann B, Buchner T, Hiddemann W. 1998. Failure of fluconazole prophylaxis to reduce mortality or the requirement of systemic amphotericin B therapy during treatment for refractory acute myeloid leukemia: results of a prospective randomized phase III study. German AML Cooperative Group. *Cancer* 83:291–301.
 70. Huijgens PC, Simoons-Smit AM, van Loenen AC, Prooy E, van Tinteren H, Ossenkoppele GJ, Jonkhoff AR. 1999. Fluconazole versus itraconazole for the prevention of fungal infections in haematology-oncology. *J Clin Pathol* 52:376–380. <https://doi.org/10.1136/jcp.52.5.376>.
 71. Kelsey SM, Goldman JM, McCann S, Newland AC, Scarffe JH, Oppenheim BA, Mufti GJ. 1999. Liposomal amphotericin (AmBisome) in the prophylaxis of fungal infections in neutropenic patients: a randomised, double-blind, placebo-controlled study. *Bone Marrow Transplant* 23:163–168. <https://doi.org/10.1038/sj.bmt.1701543>.
 72. Menichetti F, Del Favero A, Martino P, Bucaneve G, Micozzi A, Girmenia C, Barbabietola G, Pagano L, Leoni P, Specchia G, Caiozzo A, Raimondi R, Mandelli F, Cascavilla N, Rigolin G, Deliliers GL, Martino B, Peta A, Ricci P, Fanci R, De Plano W, Sivera P, Zagonel V, Caramatti C, Invernizzi R, Buelli M, Bernasconi C, Carella A, Damasio E, De Biasi R, Chierichini A, Galieni P, Grasso M, Jacopino P, Morra E, Porcellini A, Resegotti L, Ricciuti F, Picardi M, Ronconi F, Cimino R, Cimino R, Ferrandina C, Gabbas A, Mettivor V. 1999. Itraconazole oral solution as prophylaxis for fungal infections in neutropenic patients with hematologic malignancies: a randomized, placebo-controlled, double-blind, multicenter trial. *Clin Infect Dis* 28:250–255. <https://doi.org/10.1086/515129>.
 73. Morgenstern GR, Prentice AG, Prentice HG, Ropner JE, Schey SA, Warnock DW. 1999. A randomized controlled trial of itraconazole versus fluconazole for the prevention of fungal infections in patients with haematological malignancies. *Br J Haematol* 105:901–911. <https://doi.org/10.1046/j.1365-2141.1999.01465.x>.
 74. Rotstein C, Bow EJ, Laverdiere M, Ioannou S, Carr D, Moghaddam N. 1999. Randomized placebo-controlled trial of fluconazole prophylaxis for neutropenic cancer patients: benefit based on purpose and intensity of cytotoxic therapy. *Clin Infect Dis* 28:331–440. <https://doi.org/10.1086/515128>.
 75. Schwartz S, Behre G, Heinemann V, Wandt H, Schilling E, Arning M, Trittin A, Kern WV, Boenisch O, Bosse D, Lenz K, Ludwig WD, Hiddemann W, Siebert W, Beyer J. 1999. Aerosolized amphotericin B inhalations as prophylaxis of invasive aspergillus infections during prolonged neutropenia: results of a prospective randomized multicenter trial. *Blood* 93:3654–3661.
 76. Young GA, Bosly A, Gibbs DL, Durrant S. 1999. A double-blind comparison of fluconazole and nystatin in the prevention of candidiasis in

- patients with leukaemia. Antifungal Prophylaxis Study Group. *Eur J Cancer* 35:1208–1213. [https://doi.org/10.1016/S0959-8049\(99\)00102-1](https://doi.org/10.1016/S0959-8049(99)00102-1).
77. Housseau JL, Dekker AW, Stamatoullas-Bastard A, Fassas A, Linkesch W, Gouveia J, De Bock R, Rovira M, Seifert WF, Joosen H, Peeters M, De Beule K. 2000. Itraconazole oral solution for primary prophylaxis of fungal infections in patients with hematological malignancy and profound neutropenia: a randomized, double-blind, double-placebo, multicenter trial comparing itraconazole and amphotericin B. *Antimicrob Agents Chemother* 44:1887–1893. <https://doi.org/10.1128/AAC.44.7.1887-1893.2000>.
 78. Laverdière M, Rotstein C, Bow EJ, Roberts RS, Ioannou S, Carre D, Moghaddame N, Bow EJ, Evans G, Fong IW, Garber G, Haase D, Horn R, Laverdière M, Phillips P, Rotstein C, Saginur R, Salit I, Sinave C, Smaill F. 2000. Impact of fluconazole prophylaxis on fungal colonization and infection rates in neutropenic patients. *J Antimicrob Chemother* 46:1001–1008. <https://doi.org/10.1093/jac/46.6.1001>.
 79. Nucci M, Biasoli I, Akiti T, Silveira F, Solza C, Barreiros G, Spector N, Derossi A, Pulcheri W. 2000. A double-blind, randomized, placebo-controlled trial of itraconazole capsules as antifungal prophylaxis for neutropenic patients. *Clin Infect Dis* 30:300–305. <https://doi.org/10.1086/313654>.
 80. Wolff SN, Fay J, Stevens D, Herzig RH, Pohlman B, Bolwell B, Lynch J, Ericson S, Freytes CO, LeMaistre F, Collins R, Pineiro L, Greer J, Stein R, Goodman SA, Dummer S. 2000. Fluconazole vs low-dose amphotericin B for the prevention of fungal infections in patients undergoing bone marrow transplantation: a study of the North American Marrow Transplant Group. *Bone Marrow Transplant* 25:853–859. <https://doi.org/10.1038/sj.bmt.1702233>.
 81. Boogaerts M, Maertens J, Van Hoof A, De Bock R, Fillet G, Peetermans M, Selleslag D, Vandercam B, Vandewoude K, Zachée P, De Beule K. 2001. Itraconazole versus amphotericin B plus nystatin in the prophylaxis of fungal infections in neutropenic cancer patients. *J Antimicrob Chemother* 48:97–103. <https://doi.org/10.1093/jac/48.1.97>.
 82. Koh LP, Kurup A, Goh YT, Fook-Chong SM, Tan PH. 2002. Randomized trial of fluconazole versus low-dose amphotericin B in prophylaxis against fungal infections in patients undergoing hematopoietic stem cell transplantation. *Am J Hematol* 71:260–267. <https://doi.org/10.1002/ajh.10234>.
 83. Kaptan K, Uğur Ural A, Çetin T, Avcu F, Beyan C, Yalçın A. 2003. Itraconazole is not effective for the prophylaxis of fungal infections in patients with neutropenia. *J Infect Chemother* 9:40–45. <https://doi.org/10.1007/s10156-002-0207-5>.
 84. Lass-Flörl C, Gunsilius E, Gastl G, Englisch M, Koch G, Ulmer H, Dierich MP, Petzer A. 2003. Fungal colonization in neutropenic patients: a randomized study comparing itraconazole solution and amphotericin B solution. *Ann Hematol* 82:565–569. <https://doi.org/10.1007/s00277-003-0666-5>.
 85. Marr KA, Crippa F, Leisenring W, Hoyle M, Boeckh M, Balajee SA, Nichols WG, Musher B, Corey L. 2004. Itraconazole versus fluconazole for prevention of fungal infections in patients receiving allogeneic stem cell transplants. *Blood* 103:1527–1533. <https://doi.org/10.1182/blood-2003-08-2644>.
 86. Van Burik JAH, Ratanatharathorn V, Stepan DE, Miller CB, Lipton JH, Vesole DH, Bunin N, Wall DA, Hiemenz JW, Satoh Y, Lee JM, Walsh TJ. 2004. Micafungin versus fluconazole for prophylaxis against invasive fungal infections during neutropenia in patients undergoing hematopoietic stem cell transplantation. *Clin Infect Dis* 39:1407–1416. <https://doi.org/10.1086/422312>.
 87. Hiemenz J, Cagnoni P, Simpson D, Devine S, Chao N, Keirns J, Lau W, Facklam D, Buell D. 2005. Pharmacokinetic and maximum tolerated dose study of micafungin in combination with fluconazole versus fluconazole alone for prophylaxis of fungal infections in adult patients undergoing a bone marrow or peripheral stem cell transplant. *Antimicrob Agents Chemother* 49:1331–1336. <https://doi.org/10.1128/AAC.49.4.1331-1336.2005>.
 88. Glasmacher A, Cornely O, Ullmann AJ, Wedding U, Bodenstern H, Wandt H, Boewer C, Pasold R, Wolf HH, Hanel M, Dolken G, Junghans C, Andreesen R, Bertz H. 2006. An open-label randomized trial comparing itraconazole oral solution with fluconazole oral solution for primary prophylaxis of fungal infections in patients with hematological malignancy and profound neutropenia. *J Antimicrob Chemother* 57:317–325. <https://doi.org/10.1093/jac/dki440>.
 89. Mattiuzzi GN, Alvarado G, Giles FJ, Ostrosky-Zeichner L, Cortes J, O'Brien S, Verstovsek S, Faderl S, Zhou X, Raad II, Bekele BN, Leitz GJ, Lopez-Roman I, Estey EH. 2006. Open-label, randomized comparison of itraconazole versus caspofungin for prophylaxis in patients with hematologic malignancies. *Antimicrob Agents Chemother* 50:143–147. <https://doi.org/10.1128/AAC.50.1.143-147.2006>.
 90. Oren I, Rowe JM, Sprecher H, Tamir A, Benyamini N, Akria L, Gorelik A, Dally N, Zuckerman T, Haddad N, Fineman R, Dann EJ. 2006. A prospective randomized trial of itraconazole vs fluconazole for the prevention of fungal infections in patients with acute leukemia and hematopoietic stem cell transplant recipients. *Bone Marrow Transplant* 38:127–134. <https://doi.org/10.1038/sj.bmt.1705418>.
 91. Penack O, Schwartz S, Martus P, Reinwald M, Schmidt-Hieber M, Thiel E, Blau IW. 2006. Low-dose liposomal amphotericin B in the prevention of invasive fungal infections in patients with prolonged neutropenia: results from a randomized, single-center trial. *Ann Oncol* 17:1306–1312. <https://doi.org/10.1093/annonc/mdl128>.
 92. Ito Y, Ohyashiki K, Yoshida I, Takeuchi M, Aoyama Y, Mugitani A, Matsuura Y, Wakita H, Matsuda M, Sakamoto E, Kiguchi T, Urabe A, Tamura K, Kanamaru A, Masaoka T, Hasegawa H, Hino M, Hirai M, Ueda Y, Shigeno K, Ohnishi K, Iwasaki H, Goto T, Ozaki K, Iijima K, Sakai C, Miyamura K, Toubai T, Imamura M, Nakagawa Y, Suzuki K, Akiyama N, Shimada T, Mizorogi F, Wada H, Mihara H, Uoshima N, Tashima M, Taniwaki M, Miyata A, Hirokawa M, Sawada K, Watanabe K, Sugawara H, Kanakura Y, Tamaki T, Misawa M, Hara H, Karasawa H, Agou H. 2007. The prophylactic effect of itraconazole capsules and fluconazole capsules for systemic fungal infections in patients with acute myeloid leukemia and myelodysplastic syndromes: a Japanese multicenter randomized, controlled study. *Int J Hematol* 85:121–127. <https://doi.org/10.1532/IJH97.06079>.
 93. Vehreschild JJ, Bohme A, Buchheidt D, Arenz D, Harnischmacher U, Heussel CP, Ullmann AJ, Mousset S, Hummel M, Frommolt P, Wassmer G, Drziska I, Cornely OA. 2007. A double-blind trial on prophylactic voriconazole (VRC) or placebo during induction chemotherapy for acute myelogenous leukaemia (AML). *J Infect* 55:445–449. <https://doi.org/10.1016/j.jinf.2007.07.003>.
 94. Hiramatsu Y, Maeda Y, Fujii N, Saito T, Nawa Y, Hara M, Yano T, Asakura S, Sunami K, Tabayashi T, Miyata A, Matsuoka KI, Shinagawa K, Ikeda K, Matsuo K, Tanimoto M. 2008. Use of micafungin versus fluconazole for antifungal prophylaxis in neutropenic patients receiving hematopoietic stem cell transplantation. *Int J Hematol* 88:588–595. <https://doi.org/10.1007/s12185-008-0196-y>.
 95. Rijnders BJ, Cornelissen JJ, Slobbe L, Becker MJ, Doorduyn JK, Hop WC, Ruijgrok EJ, Lowenberg B, Vulto A, Lugtenburg PJ, de Marie S. 2008. Aerosolized liposomal amphotericin B for the prevention of invasive pulmonary aspergillosis during prolonged neutropenia: a randomized, placebo-controlled trial. *Clin Infect Dis* 46:1401–1408. <https://doi.org/10.1086/586739>.
 96. Sawada A, Sakata N, Higuchi B, Takeshita Y, Ishihara T, Sakata A, Kuroki M, Kondo O, Koyama M, Hirano S, Yasui M, Inoue M, Yoshioka A, Kawa K. 2009. Comparison of micafungin and fosfluconazole as prophylaxis for invasive fungal infection during neutropenia in children undergoing chemotherapy and hematopoietic stem cell transplantation. *Jpn J Clin Hematol* 50:1692–1699.
 97. Marks DI, Pagliuca A, Kibbler CC, Glasmacher A, Heussel CP, Kantecki M, Miller PJ, Ribaud P, Schlamm HT, Solano C, Cook G, IMPROVIT Study Group. 2011. Voriconazole versus itraconazole for antifungal prophylaxis following allogeneic haematopoietic stem-cell transplantation. *Br J Haematol* 155:318–327. <https://doi.org/10.1111/j.1365-2141.2011.08838.x>.
 98. Chaftari AM, Hachem RY, Ramos E, Kassis C, Campo M, Jiang Y, Prince RA, Wang W, Raad II. 2012. Comparison of posaconazole versus weekly amphotericin B lipid complex for the prevention of invasive fungal infections in hematopoietic stem-cell transplantation. *Transplantation* 94:302–308.
 99. Huang X, Chen H, Han M, Zou P, Wu D, Lai Y, Huang H, Chen X, Liu T, Zhu H, Wang J, Hu J. 2012. Multicenter, randomized, open-label study comparing the efficacy and safety of micafungin versus itraconazole for prophylaxis of invasive fungal infections in patients undergoing hematopoietic stem cell transplant. *Biol Blood Marrow Transplant* 18:1509–1516. <https://doi.org/10.1016/j.bbmt.2012.03.014>.
 100. Park HJ, Park M, Han M, Nam BH, Koh KN, Im HJ, Lee JW, Chung NG, Cho B, Kim HK, Yoo KH, Koo HH, Kang HJ, Shin HY, Ahn HS, Lim YT, Kook H, Lyu CJ, Hah JO, Park JE, Lim YJ, Seo JJ. 2014. Efficacy and safety of micafungin for the prophylaxis of invasive fungal infection during neutropenia in children and adolescents undergoing allogeneic hematopoietic SCT. *Bone Marrow Transplant* 49:1212–1216. <https://doi.org/10.1038/bmt.2014.136>.